## Photodynamic Therapy with 20% Levulanic Acid & Blue Light Skin Care After Treatment

To achieve the best results from your treatment, follow these guidelines:

- If you have a medical history of herpes simplex, please take Valtrex as directed.
- AVOID ANY SUNLIGHT AND ULTRAVIOLET EXPOSURE! YOU MUST STAY INSIDE FOR THE FIRST 48 HOURS & AVOID SUN EXPOSURE FOR TWO WEEKS FOLLOWING THIS TREATMENT.
- Prevent trauma to the treated area use a gentle facial cleanser, avoid harsh rubbing for 24 hours.
- Use Soothing Balm up to 3 times daily to reduce redness and itching.
- Use Hydrating Complex Serum 1-2 times daily for moisture and to control flaking.
- Use Gentle Vitamin-Infused Anti-Oxidant Gel to calm skin.
- You may take Motrin or Tylenol for any swelling and/or pain.
- Use ice gel packs for discomfort 20 minutes on, then 20 minutes off.
- Use Hydrating Anti-Oxidant Tinted Sunscreen, even when the sun is not out, for unavoidable sun exposure.
- Avoid Renova, Retin-A, Tretinoin, Tazorac, Retinol 20 or Retinol 50.
- Avoid hot baths or aerobic exercise for 48 hours increases burning sensation.
- Use makeup unless skin is broken.
- Treat any open or crusting skin areas with an antibiotic cream of your choice.
- DO NOT have other skin care procedures for 2 weeks following this treatment.
- Please call our office (860-243-3020) if you have any questions or concerns.

Pennoyer MD Dermatology
47 Jolley Drive
Bloomfield, CT 06002
Phone: 860-243-3020 Fax: 860-243-3002
www.drjenniferpennoyer.com